arXiv:1012.3366v3 [quant-ph] 31 Dec 2010

Two-electron entanglement in a two-dimensional isotropic harmonic trap: Radial correlation effects in the low density limit.

Przemysław Kościk, Institute of Physics, Jan Kochanowski University
ul. Świętokrzyska 15, 25-406 Kielce, Poland

Abstract

We study a two-dimensional system of two Coulombically interacting electrons in an external harmonic confining potential. More precisely, we present calculations for the singlet ground-state of the system. We explain the nature of the degeneracy in the spectrum of the reduced density matrix and provide detailed results for the dependencies of the entanglement on the interaction strength. Among other features, it is indicated that in the limit of an infinitely strong interaction, the angular and radial correlations are asymptotically independent of each other. Appearing in this limit, the pure radial correlation effects are quantitatively investigated for the first time.

Keywords: Schmidt decomposition, entanglement

1. Introduction

Studies of quantum systems trapped in confining potentials have attracted attention because of their possible applications in quantum information technology [1]. The simplest candidate for studying entanglement properties is the system composed of two Coulombically interacting electrons in an external harmonic potential, which serves well as a model of a quantum dot (QD)[2]. In this Letter we consider a two-dimensional (2D) system with a Hamiltonian

$$\hat{H} = \sum_{i=1}^{2}[\frac{\mathbf{p}_i^2}{2m_*} + \frac{\omega^2 m_*}{2} r_i^2] + \frac{e^2}{\varepsilon_*|\mathbf{r}_2 - \mathbf{r}_1|}, \tag{1}$$

where m_* is the effective electron mass and ε_* is the effective dielectric constant. This system was considered in various theoretical contexts [3, 4, 5, 6, 7] but to the best of our knowledge its entanglement properties have not been studied extensively in the literature.

Since the Hamiltonian (1) is spin independent its stationary states possess the form

$$\Psi(\zeta_1, \zeta_2) = \psi(\mathbf{r}_1, \mathbf{r}_2)\varrho(s_1, s_2), \tag{2}$$

where ϱ denotes a spin function. After the scaling $\mathbf{r} \mapsto \sqrt{\frac{2\hbar}{m^*\omega}}\mathbf{r}$, $E \mapsto \frac{\hbar\omega E}{2}$, the Schrödinger equation takes the form

$$H\psi(\mathbf{r}_1, \mathbf{r}_2) = E\psi(\mathbf{r}_1, \mathbf{r}_2), \tag{3}$$

Email address: koscik@pu.kielce.pl (Przemysław Kościk, Institute of Physics, Jan Kochanowski University ul. Świętokrzyska 15, 25-406 Kielce, Poland)

with

$$H = \sum_{i=1}^{2}[-\frac{1}{2}\triangle_{\mathbf{r}_i} + 2r_i^2] + \frac{g}{|\mathbf{r}_2 - \mathbf{r}_1|}, \tag{4}$$

where the dimensionless coupling $g = \frac{e^2}{\varepsilon^*}\sqrt{\frac{2m^*}{\omega\hbar^3}}$ represents the ratio of Coulomb interaction strength, which is related to the so-called Wigner parameter R_w of Ref. [4] by $g = \sqrt{2}R_w$.

Introducing the centre-of-mass (cm.) $\mathbf{R} = \frac{1}{2}(\mathbf{r}_1 + \mathbf{r}_2)$ and relative (rel.) $\mathbf{r} = \mathbf{r}_2 - \mathbf{r}_1$ coordinates, the Hamiltonian (4) is separated into $H = H^{\mathbf{R}} + H^{\mathbf{r}}$, where the cm. Hamiltonian $H^{\mathbf{R}} = -\nabla_R^2/4 + 4R^2$ is exactly solvable. The problem is thus reduced to the Schrödinger equation for the rel. motion

$$H^{\mathbf{r}}\psi^r = \varepsilon^{\mathbf{r}}\psi^r,$$

described by the Hamiltonian $H^{\mathbf{r}} = -\nabla_r^2 + r^2 + \frac{g}{r}$. For the sake of simplicity we restrict our investigation to the study of the singlet ground-state. The total wavefunction is given by

$$\begin{aligned}\Psi(\zeta_1, \zeta_2) &= \psi(r, R)\varrho_S(s_1, s_2) = \\ &= \frac{2}{\sqrt{\pi}}e^{-2R^2}\frac{1}{\sqrt{2\pi}}\frac{u^r(r)}{\sqrt{r}}\varrho_S(s_1, s_2),\end{aligned} \tag{5}$$

where the unknown function u^r ($\int_0^\infty [u^r]^2 dr = 1$) is the lowest-energy eigenfunction of the rel. radial equation

$$[-\frac{d^2}{dr^2} - \frac{1}{4r^2} + r^2 + \frac{g}{r}]u^r(r) = \varepsilon^{\mathbf{r}}u^r(r), \tag{6}$$

and $\varrho_S = \frac{1}{\sqrt{2}}(\alpha(1)\beta(2) - \alpha(2)\beta(1))$ is the spin singlet function, where $\alpha(\beta)$ denotes the up (down), $\uparrow$ ($\downarrow$), spin. We note that $r = \sqrt{r_1^2 + r_2^2 - 2r_1r_2cos(\varphi_2 - \varphi_1)}$ and $R = \sqrt{r_1^2 + r_2^2 + 2r_1r_2cos(\varphi_2 - \varphi_1)}/2$, so that in a correlated description ψ depends explicitly on r_1, r_2 and the inter-electronic angle coordinate $\theta = \varphi_2 - \varphi_1$.

In this Letter we will analyse the entanglement contained in the state $|\Psi\rangle$ (5). Special attention is given to performing both the partial-wave expansion and the Schmidt decomposition of ψ. We make a detailed investigation of the nature of the degeneracy in the spectrum of the reduced density matrix. We discuss the effect of interaction on various characteristics such as the collective occupancies, the participation ratio, and the Slater rank as well, over a wide range of values of g. In particular, we show that only the Schmidt modes with nodeless radial functions contribute significantly to the total wavefunction. Importantly, we give an indication that in the $g \to \infty$ ($\omega \to 0$) limit the global correlation is fully separable into purely angular and radial correlations. Moreover, in the regime of $g \to \infty$ we investigate for the first time the role played by the radial correlation on the electron pair.

The Letter is arranged as follows. In Section 2 we discuss the characteristics of the system under consideration. Section 3 is devoted to the results and a brief summary of our conclusions is provided in Section 4.

2. Entanglement characteristics

A tool to investigate two-body correlations is the reduced density matrix (RDM) defined as [8]

$$\rho_{red}(\zeta, \zeta^{'}) = tr_{\zeta_2}(|\Psi\rangle\langle\Psi|), \tag{7}$$

where tr denotes a trace taken over one of the electrons, and Ψ is the total two-electron wavefunction. As Ψ is given by Eq. (2), the RDM factors into spatial and spin components

$$\rho_{red}(\zeta,\zeta^{'}) = \rho(\mathbf{r},\mathbf{r}^{'})\rho_S(s,s^{'}), \tag{8}$$

where the spatial RDM is given by $\rho(\mathbf{r},\mathbf{r}^{'}) = \int \psi^*(\mathbf{r},\mathbf{r}_2)\psi(\mathbf{r}^{'},\mathbf{r}_2)d\mathbf{r}_2$. In particular, for the singlet states, the spin part of the RDM is a matrix $\rho_S = diag(1/2, 1/2)$. It is well known that for identical fermions the bi-partite pure state $|\Psi\rangle$ can be expressed as a combination of the Slater determinants made out of one-particle spin-orbitals in which the RDM (7) is diagonal [9]. The number of non-zero expansion coefficients in the Slater decomposition is called the Slater rank (SR) and a pure fermion state is entangled if, and only if, its Slater rank is larger than 1. The entanglement depends on the whole spectrum of the RDM and many ways of measuring its amount have been developed in the literature [10, 11, 12, 13]. It is interesting to point out that most of the popular entanglement measures are functions of the purity of the RDM, $tr\rho_{red}^2$. The best known are the participation ratio [10] and the linear entropy [11]. Here we consider the participation ratio,

$$\mathbf{R} = [tr\rho_{red}^2]^{-1}, \tag{9}$$

which approximately counts the number of Slater orbitals actively involved in the Slater expansion of Ψ. The effective Slater rank SR is defined by $SR = \mathbf{R}/2$, since the Slater determinant is made up of two different spin orbitals. The larger the value of SR, the higher the entanglement or quantum correlation. It is worth stressing at this point that the linear correlation entropy $\mathbf{L}$, which is the popular measure of entanglement in pure states [14, 15, 16, 17, 18], is related to $\mathbf{R}$ via $\mathbf{L} = 1 - 1/\mathbf{R}$.

2.1. The Slater-Schmidt decomposition

The Slater decomposition of the total wavefunction of (2) can easily be inferred from the Schmidt decomposition of the spatial part ψ. In a previous work [19], we have shown that for any 2D system the real spatial wavefunction that depends only on r and R, $\psi(r,R) \equiv \psi(r_1, r_2, cos\theta)$, can be expanded as

$$\begin{aligned}\psi(r,R) &= \sum_{m=-\infty\ldots\infty} \frac{A_m(r_1,r_2)}{\sqrt{r_1 r_2}} \frac{e^{im\varphi_1}}{\sqrt{2\pi}} \frac{e^{-im\varphi_2}}{\sqrt{2\pi}} = \\ &= \sum_{\substack{m=-\infty\ldots\infty \\ s=0,\infty}} \kappa_{s,m} u_{s,m}(\mathbf{r}_1) u^*_{s,m}(\mathbf{r}_2),\end{aligned} \tag{10}$$

where $A_m = A_{-m}$,

$$\begin{aligned}A_m(r_1,r_2) &= \frac{\sqrt{r_1 r_2}}{2\pi} \int_0^{2\pi}\int_0^{2\pi} \psi(r,R) e^{-im\varphi_1} e^{im\varphi_2} d\varphi_1 d\varphi_2 = \\ &= \sqrt{r_1 r_2} \int_0^{2\pi} \psi(r_1, r_2, cos\theta) cos(m\theta) d\theta,\end{aligned} \tag{11}$$

and

$$A_m(r_1,r_2) = \sum_{s=0}^{\infty} \kappa_{s,m} \chi_{s,m}(r_1) \chi_{s,m}(r_2), \tag{12}$$

$$u_{s,m}(\mathbf{r}) = \frac{\chi_{s,m}(r)}{\sqrt{r}} \frac{e^{im\varphi}}{\sqrt{2\pi}}. \tag{13}$$

The radial orbitals $\chi_{s,m}(r)$ and the coefficients $\kappa_{s,m}$ are determined by the integral equation

$$\int_0^\infty A_m(r_1, r_2)\chi_{s,m}(r_2)dr_2 = \kappa_{s,m}\chi_{s,m}(r_1), \tag{14}$$

and the family $\{u_{s,m}(\mathbf{r})\}$ forms a complete and orthogonal set ($\langle u_{sm}|u_{s'm'}\rangle = \int_0^\infty \int_0^{2\pi} r u^*_{sm} u_{s'm'} dr d\varphi$ $= \delta_{mm'} \int_0^\infty \chi_{sm}\chi_{s'm} dr = \delta_{mm'}\delta_{ss'}$), which means that the latter expansion appearing in Eq. (10) represents the Schmidt decomposition of ψ. Since the components A_m and A_{-m} are the same, Eq. (10) can be written in a partial form

$$\psi(r, R) = \frac{A_0(r_1, r_2)}{2\pi\sqrt{r_1 r_2}} + \sum_{l=1}^{\infty} \frac{A_l(r_1, r_2)}{\sqrt{r_1 r_2}} \frac{cos(l\theta)}{\pi}, \tag{15}$$

and $\chi_{s,m} = \chi_{s,-m}$, $\kappa_{s,m} = \kappa_{s,-m}$. The latter means that the Schmidt decomposition possesses a degenerate spectrum, and so it fails to be unique [9]. For the sake of completeness, we give below another from Eq. (10) form of the Schmidt decomposition of ψ. To begin with, we extend the results of Ref. [9] to the case with more than one point of double degeneracy. Accordingly, from the orbitals $u_{s,l}$ and $u_{s,-l}$ (13) ($l > 0$), that correspond to the same Schmidt coefficient $\kappa_{s,l}$, we define the new orbitals as

$$\nu_{s,l}(\mathbf{r}) = \frac{u_{s,l}(\mathbf{r}) + u_{s,-l}(\mathbf{r})}{\sqrt{2}}, \upsilon_{s,l}(\mathbf{r}) = i\frac{u_{s,-l}(\mathbf{r}) - u_{s,l}(\mathbf{r})}{\sqrt{2}},$$

and for $l = 0$ as $\nu_{s,0} = \frac{\chi_{s,0}(r)}{\sqrt{2\pi}\sqrt{r}} = u_{s,0}$, $\upsilon_{s,0} = 0$. The angular parts of the new Schmidt modes can be obtained analytically. After some tedious algebra, one arrives at

$$\nu_{s,l}(\mathbf{r}) = \frac{\chi_{s,l}(r)}{\sqrt{r}} \frac{cos(l\varphi)}{\sqrt{\pi}}, \upsilon_{s,l}(\mathbf{r}) = \frac{\chi_{s,l}(r)}{\sqrt{r}} \frac{sin(l\varphi)}{\sqrt{\pi}}. \tag{16}$$

The family $\{\nu_{s,l}, \upsilon_{s,l}\}$ forms a complete and orthonormal set, since one can easily check that $\langle \nu_{sl}|\nu_{s'l'}\rangle = \delta_{ss'}\delta_{ll'}$, $\langle \upsilon_{s,l}|\upsilon_{s'l'}\rangle = \delta_{ss'}\delta_{ll'}$, $\langle \nu_{s,l}|\upsilon_{s'l'}\rangle = 0$. In terms of the new orbitals, the wave-function takes the form

$$\psi(r, R) = \sum_{l,s=0}^{\infty} \kappa_{s,l}[\nu_{s,l}(\mathbf{r}_1)\nu_{s,l}(\mathbf{r}_2) + \upsilon_{s,l}(\mathbf{r}_1)\upsilon_{s,l}(\mathbf{r}_2)], \tag{17}$$

which yields a Schmidt form different from Eq. (10). Consequently, there exist also two different forms of the Slater expansion of (5), which can easily be inferred from the Schmidt decompositions (10) and (17).

2.2. The partial expansion of the RDM

Now using Eq. (10) we turn the spatial RDM into the partial form

$$\rho(\mathbf{r}, \mathbf{r}') = \frac{\rho_0(r, r')}{2\pi\sqrt{rr'}} + \sum_{l=1}^{\infty} \frac{\rho_l(r, r')}{\sqrt{rr'}} \frac{cos(l\theta')}{\pi}, \tag{18}$$

wherein $\theta^{'} = \varphi - \varphi^{'}$, and ρ_l (the l-matrix) is given by

$$\rho_l(r, r^{'}) = \int_0^\infty A_l(r, r_2) A_l(r^{'}, r_2) dr_2. \tag{19}$$

It can easily be derived that the l-matrix has the expansion

$$\rho_l(r, r^{'}) = \sum_{s=0}^{\infty} \lambda_{s,l} \chi_{s,l}(r) \chi_{s,l}(r^{'}), \tag{20}$$

which represents nothing else but its Schmidt decomposition. The eigenequation of the l-matrix is thus of the form

$$\int_0^\infty \rho_l(r, r^{'}) \chi_{s,l}(r^{'}) dr^{'} = \lambda_{s,l} \chi_{s,l}(r). \tag{21}$$

Obviously, the spatial RDM (18) is diagonal in the basis of Schmidt modes (spatial natural orbitals) and its eigenvalues $\lambda_{s,l}$ (occupancies) are related to $\kappa_{s,l}$ via the formula $\lambda_{s,l} = \kappa_{s,l}^2$. Moreover, all occupancies but the ones with $l = 0$ are doubly degenerate so that the normalization condition gives $\sum_{s=0}^{\infty} \lambda_{s,0} + 2 \sum_{l=1}^{\infty} \sum_{s=0}^{\infty} \lambda_{s,l} = 1$.

The quantity of interest is the sum of the occupancies corresponding to the l-matrix $\eta_l = \sum_{s=0}^{\infty} \lambda_{s,l}$. Throughout this Letter we will refer to $\{\eta_l\}$ as the collective occupancies. They can simply be calculated by $\eta_l = Tr\rho_l = \int_0^\infty \rho_l(r, r) dr = ||A_l||^2$, which is crucial since for their determination the numerical diagonalizations of the matrices ρ_l are not needed. It is clear that, if η_l is vanishingly small compared with the remaining collective occupancies, we may expect that the l-partial wave in Eq. (15) contributes little. Moreover, η_0 might be seen as the fraction of electrons with zero angular momentum, and $2\eta_l$ ($l > 0$), as the fraction of electrons with fixed opposite angular momenta ($\hbar l$ and $-\hbar l$). This is particularly easy to understand when rewriting the spatial wavefunction as a combination of symmetrized products of the modes $u_{s,l}$, $u_{s,-l}$ (13) [1].

2.3. The participation ratio

The purity of the RDM, $tr\rho_{red}^2$, of the singlet state separates in $\frac{1}{2} tr\rho^2$, where $\frac{1}{2}$ is related to the spin degree of freedom. Moreover, in the case of (18), $tr\rho^2$ can be decomposed as

$$tr\rho^2 = tr\rho_0^2 + 2 \sum_{l=1}^{\infty} tr\rho_l^2, \tag{22}$$

where

$$tr\rho_l^2 = \int_0^\infty \rho_l^2(r, r) dr, \tag{23}$$

with

$$\rho_l^2(r_1, r_2) = \int_0^\infty \rho_l(r_1, r_3) \rho_l(r_3, r_2) dr_3. \tag{24}$$

Accordingly, in terms of (22) the participation ratio of the singlet state can be written as

$$\mathbf{R} = [\frac{1}{2} tr\rho_0^2 + \sum_{l=1}^{\infty} tr\rho_l^2]^{-1}. \tag{25}$$

[1] $\psi(r, R) = \sum_{s=0}^{\infty} \kappa_{s,0} u_{s,0}(\mathbf{r}_1) u_{s,0}(\mathbf{r}_2) + \sum_{s=0,l=1}^{\infty} \sqrt{2} \kappa_{s,l} [\frac{1}{\sqrt{2}} (u_{s,l}(\mathbf{r}_1) u_{s,-l}(\mathbf{r}_2) + u_{s,l}(\mathbf{r}_2) u_{s,-l}(\mathbf{r}_1))]$

From a practical point of view the above expansion appears to be a convenient tool to determine the number N_l of partial waves that are needed to capture most of the electrons correlation. It seems to be reasonable to define N_l as the minimal number of terms in the sum in (25) at which an approximate value for $\mathbf{R}$ has its integer part equal to the integer part of the exact value of $\mathbf{R}$.

Clearly, the effective Schmidt number of non-zero coefficients in the Schmidt decomposition of A_l, Eq. (12), is determined by the participation ratio of the normalized l-matrix,

$$\bar{\rho}_l(r, r') = \frac{\rho_l(r, r')}{Tr\rho_l}, \tag{26}$$

$(Tr\bar{\rho}_l = 1)$, i.e. $[tr\bar{\rho}_l^2]^{-1}$.

3. Numerical results

To start with our analysis, we need the solutions of (6). As was shown by Taut [3] for a countably infinite set of g values the closed-form wavefuntions can be derived. $g = \sqrt{2}$ is the smallest value at which an exact ground-state wavefunction is known. Below this value no analytical solutions to the lowest eigenvalue of (6) are known and we have determined them numerically by the Rayleigh-Ritz method. As was already discussed, a qualitative insight into the character of the partial-wave expansion of ψ can be provided by examining the values of the collective occupancies. In Fig. 1 the numerically determined behaviour of η_l is presented for $l = 0 - 5$, where to highlight the changes with g a logarithmic coordinate has been used. We may notice that at $g \lesssim 1$ the collective




Figure 1: The dependence of the six lowest collective occupancies η_l $(l = 0 - 5)$ of the lowest singlet state on $\ln g$. The inset highlights their changes for large values of g.

occupancy with $l = 0$ is about 1. In that situation the angular correlation effects may be neglected since one can expect that the terms higher than $l = 0$ in (15) contribute very little. When g increases, more and more collective occupancies with different l become substantial, which reflects

the fact that the larger is the value of g, the larger is the number of partial waves actively involved in the sum in (15). Interestingly enough, all the collective occupancies *except* for the largest one (η_0) exhibit a local maximum. The larger is l, the larger is the value of g at which the maximum occurs. For large enough values of g the situation changes essentially, namely the clustering of collective occupancies is visible. It may have a general interest to note at this point that at $g \lesssim 7.4$ ($\ln 7.4 \approx 2$) the collective occupancies with $l > 1$ are vanishingly small, whereas a Wigner molecule is expected to occur when g exceeds a value $g_{cr} \simeq 2.8$ (See Fig. 5 in Ref. [6], remember that $g = R_w\sqrt{2}$). Thus, in the incipient Wigner molecule regime, the angular correlations come mainly from the first angular level ($l = 1$). Further, in the range of g in which $||A_l||^2 = \eta_l$ is substantial, we analyse the dependence of the effective Schmidt number for A_l, Eq. (12), on g. Our numerical results are presented in Fig. 2 in terms of the inverse participation ratio, $tr\bar{\rho_l}^2 = \Omega_l$, for $l = 0 - 3$ as a function of $\ln g$. The results obtained indicate nothing but that those partial waves that contribute considerably to ψ have their radial parts A_l being almost uncorrelated functions. That is, $A_l(r_1, r_2) \sim \chi_{0,l}(r_1)\chi_{0,l}(r_2)$, as we have verified by solving Eq. (14) through a discretization technique. In other words, we have the result that only the Schmidt orbitals with $(s = 0, l)$ in (17) are important and, in consequence, the total wavefunction approaches the form

$$\Psi(\zeta_1, \zeta_2) \approx \frac{1}{\sqrt{2}}(\alpha(1)\beta(2) - \alpha(2)\beta(1)) \times \sum_{l=0}^{N_l-1} \kappa_{0,l}[\nu_{0,l}(\mathbf{r}_1)\nu_{0,l}(\mathbf{r}_2) + \upsilon_{0,l}(\mathbf{r}_1)\upsilon_{0,l}(\mathbf{r}_2)], \tag{27}$$

where N_l is the number of substantial partial waves. The inspection of (27) yields that it constitutes a sum of $2N_l - 1$ Slater determinants (here we recall that $\upsilon_{0,0} = 0$). In the limit of $g \to \infty$, the values of $\{\Omega_l\}_{l=0}$ saturate to the same value $\Omega^{(\infty)} \simeq 0.9634$, which reflects the fact that at $g \to \infty$ the occupancies are asymptotically independent of the angular momentum i.e., $\lambda_{s,0}^{g\to\infty} = \lambda_{s,1}^{g\to\infty} = ... = \lambda_s^{(\infty)}$ ($\rho_0^{g\to\infty} = \rho_1^{g\to\infty} = ... = \rho^{(\infty)}$). We find this consistent with the properties of the 3D Hooke's atom that were discussed in [20]. What follows further from the above is that in the limit $g \to \infty$, the radial components $\{A_l\}_{l=0}$ can differ from each other only by their signs. We have verified that $A_l^{g\to\infty} = (-1)^l A^{(\infty)}$. Substitution of this into Eq. (15) gives

$$\psi^{g\to\infty}(r, R) = \frac{A^{(\infty)}(r_1, r_2)}{\sqrt{r_1 r_2}}[\frac{1}{2\pi} + \frac{1}{\pi}\sum_{l=1}^{\infty}(-1)^l cos(l\theta)]. \tag{28}$$

Because $(-1)^l cos(l\theta) = cos[l(\theta - \pi)]$, the suitable part of Eq. (28) can be recognized as a Dirac delta distribution centred at π i.e. $\delta(\theta - \pi)$[2]. The angular correlations carry thus the electrons strictly at opposite sides of the centre of the trap as $g \to \infty$, which corresponds to a perfect linear Wigner molecule. In this limit the potential energy dominates over the electron kinetic energy and the electrons localize to the classical positions $r_{1,2}^{cl} = \frac{1}{2}(\frac{g}{2})^{\frac{1}{3}}$, $\theta = \pi$, but always there exist quantum fluctuations with respect to their radial coordinates. The electrons oscillate about the equilibrium geometry in accordance with a harmonic approximation model [3, 5, 6]. At $g \to \infty$ there is no fluctuation in θ, but the angular correlations are extremely strong due to an arbitrary orientation of the molecular axis, which coincides with the fact that the lowest-energy classical configuration is

[2] A Fourier series expansion of $\delta(x - s)$, gives $\delta(x - s) = \frac{1}{2\pi} + \frac{1}{\pi}\sum_{l=1}^{\infty} cos[x(l - s)]$[22]

infinitely degenerate with respect to rotations around the center of the trap. When it comes to pure radial correlations appearing in the asymptotic regime they are expected to be very poor, due to the closeness of the value of $\Omega^{(\infty)}$ to unity ($\Omega^{(\infty)} \simeq 0.9634$). More precisely, we have found numerically that the ratio of $\lambda_0^{(\infty)}/\lambda_s^{(\infty)}$ is about 54, 2883, 154804 for $s = 1, 2, 3$, respectively, and it tends to rapidly increase with further increasing s. In other words, the radial component appearing in Eq. (28) can indeed be considered as almost a product function, which in turn implies low pure radial correlation effects. We stress that the limiting values of the considered quantities have been obtained with the use of the wavefunction in the harmonic approximation, which is valid as $g \to \infty$. Before going further we want to stress that in none of the well known papers concerning the system under consideration has Eq. (28) appeared. To gain further understanding we next quantify the degree of entanglement by calculating the effective Slater rank $SR = \mathbf{R}/2$. An approximate value for $\mathbf{R}$ can be obtained with the help of Eq. (25) by successively increasing the number of terms in the sum appearing in it until the result becomes stable to a desired accuracy. In Fig. 3 the behaviour of SR determined by the procedure described above is shown. We have checked that




Figure 2: The behaviour of Ω_l for $l = 0 - 3$ as a function of $\ln g$. The first vertical line (on the left) corresponds to the value of g below which the partial wave with $l = 1$ hardly contributes to ψ at all. The second line concerns the same but for $l = 2$, and so on. The locations of the lines were roughly inferred from Fig. 1.

the results for the linear entropy calculated with Eq. (25) as $\mathbf{L} = 1 - 1/\mathbf{R}$ completely agree with the results presented in Ref. [21], where they were calculated in a different way. The circles in Fig. 3 represent the values of the Slater rank obtained from our earlier finding SR$= 2N_l - 1$ with N_l being determined as was discussed at the end of subsection 2.3. As one can see, the behaviour of SR coincides fairy well with that of SR, which justifies the approximation (27).

4. Summary

In conclusion, we have investigated the nature of the degeneracy in the spectrum of the spatial RDM of the two-particle wavefunction that is a function of distances r and R only. We have developed the results of Ref. [19, 23] by providing the Schmidt decomposition in the single real particle




Figure 3: The effective Slater rank $SR = \mathbf{R}/2$ as a function of $\ln g$. The circles mark the results obtained by SR$= 2N_l - 1$.

basis in parabolic coordinates. In addition, we carried out a comprehensive study of the singlet ground-state of two interacting electrons confined in an isotropic harmonic potential. As a general trend we found that as g increases, the number of partial waves that are needed to capture most of the electrons' correlation increases as well. The Slater rank grows monotonically with increasing g and the bulk of the entanglement is mainly manifested in the angular variables. At $g \to \infty$ the spatial wavefunction factorizes asymptotically into a product of radial and angular components, of which the former is individually little correlated, but the latter is maximally correlated. Our calculations have shown that only at $g \lesssim 1$ can the total wavefunction be well approximated by a single Slater determinant. In other words, in this regime the state under consideration can be regarded as weakly entangled.

It would be interesting to apply the tools presented in this Letter to explore the entanglement properties of two-particle systems with other interactions between the particles. This will be the topic of our further studies.

References

[1] R. Zak, B. Roöthlisberger, S. Chesi, and D. Loss, Lecture notes for Course CLXXI "Quantum Coherence in Solid State Systems" Int. School of Physics "Enrico Fermi," Varenna, July 2008, cond-mat/0906.4045.
[2] L. Jacak, P. Hawrylak, and A. Wójs, Quantum Dots, Springer, Berlin, 1997.
[3] M. Taut, J. Phys. A: Math. Gen. 27, 1045 (1994).
[4] C. Yannouleas and U. Landman, Phys. Rev. Lett. 85, 1726 (2000).
[5] A. Matulis and F. Peeters, Solid State Commun. 117, 655 (2001).
[6] A. Puente, L. Serra, and R. Nazmitindov, Phys. Rev. B 69, 125315 (2004)
[7] H. Hassanabadia, Eur. Phys. J. B 74, 415–418 (2010)
[8] A. Coleman and V. Yukalov, Reduced Density Matrices, Springer-Verlag, Berlin, 2000.
[9] G. Ghirardi and L. Marinatto, Phys. Rev. A 70, 012109 (2004)
[10] R. Grobe, K. Rzążewski, and J. Eberly, J. Phys. B 27, L503 (1994).
[11] G. Manfredi and M. Feix, Phys. Rev. E 62, 4665 (2000).

[12] J. Batle et al., Phys. Rev. A 024301 (2005)
[13] R. Paškauskas and L. You, Phys. Rev. A 64, 042310 (2001)
[14] F. Buscemi, P. Bordone, and A. Bertoni, Phys. Rev. A 75, 032301 (2007).
[15] J. P. Coe, A. Sudbery, and I. D'Amico, Phys. Rev. B 77, 205122 (2008).
[16] D. Manzano, A. Plastino, J. Dehesa and T. Koga,J. Phys. A: Math. Theor. 43 (2010) 275301 (14pp)
[17] R. Yañez, A. Plastino, and J. Dehesa, Eur. Phys. J. D56, 141–150 (2010).
[18] F. Carlier, A. Mandilara and A. Sarfati,J. Phys. B: At. Mol. Opt. Phys. 40 (2007) S199S207
[19] P. Kościk, Phys. Lett. A 374 2999–3000 (2010).
[20] J. Ciosłowski and M. Buchowiecki, J. Chem. Phys. 125, 064105 (2006).
[21] P. Kościk and A. Okopińska, Phys. Lett. A 374, 3841–3846 (2010).
[22] R. Bracewell, The Impulse Symbol. Ch. 5 in: The Fourier Transform and Its Applications, 3rd ed.: McGraw-Hill, New York, 1999.
[23] B. Sun, M. Pindzola Phys. Lett. A 373, 3833–3837 (2009).